Drought Information Statement for Deep South Texas Valid March 15, 2025 Issued By: NWS Brownsville/Rio Grande Valley, TX Contact Information: <u>sr-bro.webmaster@noaa.gov</u>


- This product will be updated around March 28, 2025, or sooner if drought conditions change significantly
- Please see all currently available products at <u>https://drought.gov/drought-information-statements</u>
- Please visit <u>https://www.weather.gov/bro/DroughtInformationStatement</u> for previous statements
- Please visit <u>https://www.drought.gov/drought-status-updates</u> for regional drought status updates
- Severe Drought Conditions Have Expanded Across most of Deep South Texas, including into the upper Rio Grande Valley
- Moderate Drought and Abnormally Dry conditions continue for most of the remainder of Deep South Texas, including portions of the mid to lower Rio Grande Valley

U.S. Drought Monitor

Latest U.S. Drought Monitor | Latest Drought Monitor for Deep South Texas

Drought Intensity and Extent

- Severe Drought (D2) conditions are being observed across more than 64% of Deep South Texas, including all of Zapata, Jim Hogg, Brooks, Starr, western Hidalgo, and western Kenedy counties.
- **Moderate Drought (D1)** conditions are being observed across 23% of Deep South Texas, including the remainder of Hidalgo, most of Kenedy, western Willacy, and western Cameron counties.
- Abnormally Dry (D0) conditions continue across 5% of Deep South Texas, including the remainder of Kenedy, central Willacy, and west-central Cameron counties.






Recent Change in Drought Intensity

Latest U.S. Drought Monitor Class Change | Latest 4 Week Change Map for Deep South Texas

Four Week Drought Monitor Class Change

- Drought conditions have worsened across most of Deep South Texas, including the Rio Grande Valley.
- In the past 4 weeks, there has been a 1 class degradation across Zapata, Jim Hogg, Brooks, Starr, western Hidalgo, most of Kenedy, most of Willacy, and western Cameron counties.
- There has been **no change** in drought condition across the central Kenedy and eastern Hidalgo counties.






Precipitation

National Water Prediction Services

- Very minimal rainfall has been observed across Deep South Texas over the past month, with multiple dry frontal passages.
- All of Deep South Texas has received 25% or less of normal rainfall over the past 30 days.
- Over the past 90 days, most of Deep South Texas has received between 25 to 50% of normal rainfall, with most of Zapata, Starr, southern Jim Hogg, southern Brooks, and western Hidalgo counties receiving 25% or less of normal rainfall.



National Weather Service Brownsville/Rio Grande Valley, TX


Last Updated: 03/14/25




Temperature

Daily Climate Summary: <u>BRO</u> | <u>HRL</u> | <u>MFE</u>

- <u>Average Maximum Temperatures</u> over the past 30 days across Deep South Texas have ranged generally **near to slightly above normal** between 70-80 degrees.
- <u>Average Minimum Temperatures</u> over the past 30 days across Deep South Texas have ranged generally **near to slightly below normal** between 50-60 degrees.
- Overall, above normal highs and lows are expected through Saturday, March 22, 2025, with slightly below normal to normal temperatures Thursday and Friday, March 20-21.









View or Submit: Condition Monitoring Observer Reports (CMOR) | Drought Impacts Reporter

Hydrologic Impacts

- Streamflows have remained well below normal due to the lack of beneficial rain across most basins in Deep South Texas.
- Texas water share levels at both Amistad and Falcon Lake remain low.

Agricultural Impacts

- Please see the latest <u>Crop and Weather Report</u> from Texas A&M AgriLife.
- Soil moistures generally range from near normal along the coast and lower to mid Rio Grande Valley to below normal across the brush country, with crop moisture indices near normal to abnormally dry across most of Deep South Texas.

Fire Hazard Impacts

- Above normal wildland fire activity is expected through April for most of Deep South Texas, with near normal activity May through June.
- Burn bans are in effect for all counties of Deep South Texas except Cameron County.

Mitigation Actions

<u>TCEQ Known Municipality Restrictions</u>



National Oceanic and Atmospheric Administration U.S. Department of Commerce


Hydrologic Conditions and Impacts

Current Amistad Reservoir Data | Current Falcon Lake Reservoir Data

- Streamflows over the past 7 days have remained much below normal with a continued lack of beneficial rainfall, above normal high temperatures, and dry frontal passages.
- Most of the streamflow across Deep South Texas is less than the 10th percentile for this time of year (dark red to maroon shading on the map).
- Texas water share values have remained near 25% at Amistad and near 15% at Falcon Lake.

Pool Elevation* (ft)	Current Elevation* (ft)	Percent Full*
1117.00	1051.06	26.1%
301.10	255.71	15.7%
1 Month Ago	3 Months Ago	1 Year Ago
26.3%	26.5%	28.2%
15.2%	12.8%	15.8%
	Elevation* (ft) 1117.00 301.10 1 Month Ago 26.3%	Elevation* (ft) Elevation* (ft) 1117.00 1051.06 301.10 255.71 1 Month Ago 3 Months Ago 26.3% 26.5%





Captions: Left: <u>TWDB Reservoir</u> conditions as of March 15, 2025 Right: <u>USGS 7 Day Streamflows for Texas</u> valid March 14, 2025

National Weather Service Brownsville/Rio Grande Valley, TX





National Oceanic and Atmospheric Administration U.S. Department of Commerce



Latest Crop and Weather Report from Texas A&M AgriLife | Climate Prediction Center (CPC) Drought Page

- Soil moistures range from near normal along the coast and lower to mid Rio Grande Valley to below normal towards the brush country.
- Crop moisture indices are generally near normal to abnormally dry across most of Deep South Texas.








National Oceanic and Atmospheric Administration

Fire Hazard Impacts

National Interagency Coordination Center (NICC) Wildfire Potential Outlooks

- Keetch-Byram Drought Index values generally range between 500-700 across most of Deep South Texas, with KBDI values between 300-500 across Cameron and Willacy counties.
- Above normal wildland fire potential is expected through April 2025 for most of Deep South Texas, with near normal wildland fire potential between May and June 2025.
- <u>Burn bans</u> remain in effect for all of Deep South Texas except Cameron County.






National Oceanic and Atmospheric Administration U.S. Department of Commerce

Seven Day Precipitation Forecast

CPC 6-10 Day Precipitation Outlook | WPC Precipitation Forecasts

- Little to no rainfall is expected through Saturday, March 22, 2025 across Deep South Texas.
- Overall, rain chances through Monday, March 24th, are leaning toward **below normal** across Deep South Texas.

7-Day Quantitative Precipitation Forecast for March 14, 2025–March 21, 2025







Long-Range Outlooks

CPC Seasonal Temperature Outlook | CPC Seasonal Precipitation Outlook

- There is a **50-60% probability of above normal temperatures** across Deep South Texas through the month of March.
- There is a **50-60% probability of below normal rainfall** across Deep South Texas through the month of March.
- Through May 2025, there is a likely chance of above normal temperatures and chances are leaning toward below normal rainfall across Deep South Texas.









Climate Prediction Center | Monthly Drought Outlook | Seasonal Drought Outlook

- **Drought is expected to persist** across most of Deep South Texas, including the northern ranchlands, brush country, and mid to upper Rio Grande Valley, through May 2025.
- **Drought is expected to develop** across the remainder of Deep South Texas, including coastal counties and lower Rio Grande Valley through May 2025.

Seasonal (3-Month) Drought Outlook for February 28, 2025-May 31, 2025

