

NOAA Technical Memorandum
NWS ER-102



A SEVERE WEATHER CLIMATOLOGY FOR THE WILMINGTON, OH WFO COUNTY WARNING AREA (1950-2004)

MICHAEL D. RYAN

National Weather Service Weather Forecast Office
WILMINGTON, OH

Scientific Services Division
Eastern Region Headquarters
Bohemia, New York
May 2008

NOAA TECHNICAL MEMORANDA
National Weather Service, Eastern Region Subseries

The National Weather Service Eastern Region (ER) Subseries provides an informal medium for the documentation and quick dissemination of results not appropriate, or not yet ready for formal publications. The series is used to report on work in progress, to describe technical procedures and practices, or to relate progress to a limited audience. These Technical Memoranda will report on investigations devoted primarily to regional and local problems of interest mainly to ER personnel, and usually will not be widely distributed.

Papers 1 to 22 are in the former series, ESSA Technical Memoranda, Eastern Region Technical Memoranda (ERTM); papers 23 to 37 are in the former series, ESSA Technical Memoranda, Weather Bureau Technical Memoranda (WBTM). Beginning with 38, the papers are now part of the series, NOAA Technical Memoranda NWS.

Papers 1 to 22 are available from the National Weather Service Eastern Region, Scientific Services Division, 630 Johnson Avenue, Bohemia, NY, 11716. Beginning with 23, the papers are available from the National Technical Information Service, U.S. Department of Commerce, Sills Bldg., 5285 Port Royal Road, Springfield, VA 22161. Prices vary for paper copy and for microfiche. Order by accession number shown in parentheses at end of each entry.

ESSA Technical Memoranda

| | | |
|------------|----|--|
| ERTM | 1 | Local Uses of Vorticity Prognoses in Weather Prediction. Carlos R. Dunn. April 1965. |
| ERTM | 2 | Application of the Barotropic Vorticity Prognostic Field to the Surface Forecast Problem. Silvio G. Simplicio. July 1965. |
| ERTM | 3 | A Technique for Deriving an Objective Precipitation Forecast Scheme for Columbus, Ohio. Robert Kuessner. September 1965. |
| ERTM | 4 | Stepwise Procedures for Developing Objective Aids for Forecasting the Probability of Precipitation. Carlos R. Dunn. November 1965. |
| ERTM | 5 | A Comparative Verification of 300 mb. Winds and Temperatures Based on NMC Computer Products Before and After Manual Processing. Silvio G. Simplicio. March 1966. |
| ERTM | 6 | Evaluation of OFDEV Technical Note No. 17. Richard M. DeAngelis. March 1966. |
| ERTM | 7 | Verification of Probability of Forecasts at Hartford, Connecticut, for the Period 1963-1965. Robert B. Wassall. March 1966. |
| ERTM | 8 | Forest-Fire Pollution Episode in West Virginia, November 8-12, 1964. Robert O. Weedfall. April 1966. |
| ERTM | 9 | The Utilization of Radar in Meso-Scale Synoptic Analysis and Forecasting. Jerry D. Hill. March 1966. |
| ERTM | 10 | Preliminary Evaluation of Probability of Precipitation Experiment. Carlos R. Dunn. May 1966. |
| ERTM | 11 | Final Report. A Comparative Verification of 300 mb. Winds and Temperatures Based on NMC Computer Products Before and After Manual Processing. Silvio G. Simplicio. May 1966. |
| ERTM | 12 | Summary of Scientific Services Division Development Work in Sub-Synoptic Scale Analysis and Prediction - Fiscal Year 1966. Fred L. Zuckerberg. May 1966. |
| ERTM | 13 | A Survey of the Role of Non-Adiabatic Heating and Cooling in Relation of the Development of Mid-Latitude Synoptic Systems. Constantine Zois. July 1966. |
| ERTM | 14 | The Forecasting of Extratropical Onshore Gales at the Virginia Capes. Glen V. Sachse. August 1966. |
| ERTM | 15 | Solar Radiation and Clover Temperatures. Alex J. Kish. September 1966. |
| ERTM | 16 | The Effects of Dams, Reservoirs and Levees on River Forecasting. Richard M. Greening. September 1966. |
| ERTM | 17 | Use of Reflectivity Measurements and Reflectivity Profiles for Determining Severe Storms. Robert E. Hamilton. October 1966. |
| ERTM | 18 | Procedure for Developing a Nomograph for Use in Forecasting Phenological Events from Growing Degree Days. John C. Purvis and Milton Brown. December 1966. |
| ERTM | 19 | Snowfall Statistics for Williamsport, Pa. Jack Hummel. January 1967. |
| ERTM | 20 | Forecasting Maturity Date of Snap Beans in South Carolina. Alex J. Kish. March 1967. |
| ERTM | 21 | New England Coastal Fog. Richard Fay. April 1967. |
| ERTM | 22 | Rainfall Probability at Five Stations Near Pickens, South Carolina, 1957-1963. John C. Purvis. April 1967. |
| WBTM ER 23 | | A Study of the Effect of Sea Surface Temperature on the Areal Distribution of Radar Detected Precipitation Over the South Carolina Coastal Waters. Edward Paquet. June 1967. (PB-180-612). |
| WBTM ER 24 | | An Example of Radar as a Tool in Forecasting Tidal Flooding. Edward P. Johnson. August 1967. (PB-180-613). |
| WBTM ER 25 | | Average Mixing Depths and Transport Wind Speeds over Eastern United States in 1965. Marvin E. Miller. August 1967. (PB-180-614). |
| WBTM ER 26 | | The Sleet Bright Band. Donald Marier. October 1967. (PB-180-615). |
| WBTM ER 27 | | A Study of Areas of Maximum Echo Tops in the Washington, D.C. Area During the Spring and Fall Months. Marie D. Fellechner. April 1968. (PB-179-339). |
| WBTM ER 28 | | Washington Metropolitan Area Precipitation and Temperature Patterns. C.A. Woollum and N.L. Canfield. June 1968. (PB-179-340). |
| WBTM ER 29 | | Climatological Regime of Rainfall Associated with Hurricanes after Landfall. Robert W. Schoner. June 1968. (PB-179-341). |
| WBTM ER 30 | | Monthly Precipitation - Amount Probabilities for Selected Stations in Virginia. M.H. Bailey. June 1968. (PB-179-342). |
| WBTM ER 31 | | A Study of the Areal Distribution of Radar Detected Precipitation at Charleston, S.C. S.K. Parrish and M.A. Lopez. October 1968. (PB-180-480). |
| WBTM ER 32 | | The Meteorological and Hydrological Aspects of the May 1968 New Jersey Floods. Albert S. Kachic and William Long. February 1969. (Revised July 1970). (PB-194-222). |
| WBTM ER 33 | | A Climatology of Weather that Affects Prescribed Burning Operations at Columbia, South Carolina. S.E. Wasserman and J.D. Kanupp. December 1968. (COM-71-00194). |
| WBTM ER 34 | | A Review of Use of Radar in Detection of Tornadoes and Hail. R.E. Hamilton. December 1969. (PB-188-315). |
| WBTM ER 35 | | Objective Forecasts of Precipitation Using PE Model Output. Stanley E. Wasserman. July 1970. (PB-193-378). |
| WBTM ER 36 | | Summary of Radar Echoes in 1967 Near Buffalo, N.Y. Richard K. Sheffield. September 1970. (COM-71-00310). |
| WBTM ER 37 | | Objective Mesoscale Temperature Forecasts. Joseph P. Sobel. September 1970. (COM-71-0074). |

NOAA Technical Memoranda NWS

| | | |
|-----|-------|---|
| NWS | ER 38 | Use of Primitive Equation Model Output to Forecast Winter Precipitation in the Northeast Coastal Sections of the United States. Stanley E. Wasserman and Harvey Rosenblum. December 1970. (COM-71-00138). |
| NWS | ER 39 | A Preliminary Climatology of Air Quality in Ohio. Marvin E. Miller. January 1971. (COM-71-00204). |
| NWS | ER 40 | Use of Detailed Radar Intensity Data in Mesoscale Surface Analysis. Robert E. Hamilton. March 1971. (COM-71-00573). |
| NWS | ER 41 | A Relationship Between Snow Accumulation and Snow Intensity as Determined from Visibility. Stanley E. Wasserman and Daniel J. Monte. (COM-71-00763). January 1971. |
| NWS | ER 42 | A Case Study of Radar Determined Rainfall as Compared to Rain Gage Measurements. Martin Ross. July 1971. (COM-71-00897). |
| NWS | ER 43 | Snow Squalls in the Lee of Lake Erie and Lake Ontario. Jerry D. Hill. August 1971. (COM-72-00959). |
| NWS | ER 44 | Forecasting Precipitation Type at Greer, South Carolina. John C. Purvis. December 1971. (COM-72-10332). |
| NWS | ER 45 | Forecasting Type of Precipitation. Stanley E. Wasserman. January 1972. (COM-72-10316). |

(CONTINUED ON INSIDE REAR COVER)



NOAA Technical Memorandum NWS ER-102

**A SEVERE WEATHER CLIMATOLOGY FOR THE
WILMINGTON, OH WFO COUNTY
WARNING AREA (1950-2004)**

MICHAEL D. RYAN

**National Weather Service Office
WILMINGTON, OH**

**Scientific Services Division
Eastern Region Headquarters
Bohemia, New York
May 2008**

United States
Department of Commerce
Carlos M. Gutierrez
Secretary

National Oceanic and
Atmospheric Administration
Conrad C. Lautenbaucher, Jr.
Under Secretary and Administrator

National Weather Service
John L. Hayes
Assistant Administrator

Table of Contents

| | |
|--|---|
| 1. Introduction | 1 |
| 2. Data and Methodology | 1 |
| 3. County Warning Area Topography and Demographics | 2 |
| 3.1 Topography | 2 |
| 3.2 Demographics | 2 |
| 4. Severe Weather Climatology | 2 |
| 4.1 Tornado Climatology | 3 |
| 4.2 Convective Wind Climatology | 4 |
| 4.3 Hail Climatology | 5 |
| 5. Summary and Conclusions | 5 |
| 6. Acknowledgements | 6 |
| 7. References | 6 |
| 8. Figures | 9 |

1. INTRODUCTION

The National Weather Service Forecast Office (WFO) in Wilmington, Ohio, has forecast and warning responsibilities for 52 counties across southwest Ohio, north-central Kentucky, and southeast Indiana ([Fig. 1](#)). Severe convective storms can occur at any time of the year in this area, although they are most frequent during the spring and summer.

The motivation for producing a severe weather climatology for the Wilmington, Ohio, county warning area (CWA) is to provide forecasters with a historical baseline of the likelihood and type of severe weather on annual, seasonal, monthly and hourly scales. A severe weather climatology can supply forecasters with an extensive background in assessing the frequency and magnitude of different types of severe weather on temporal and spatial scales, and assist in anticipation of local severe weather. The period of the study extends from the early 1950s to 2004.

Section 2 will explain the methodology employed in producing the data for this study. Section 3 will introduce the demographics and topography of the Wilmington CWA. Section 4 will present an analysis of the tornado, severe convective wind and hail events across the Wilmington CWA over the period of study, and provide a climatological basis for observed trends in severe weather across the Wilmington CWA. Section 5 will offer a summary of the findings. It is hoped that this study may contribute to additional climate-related research on severe convective storms affecting the Wilmington CWA.

2. DATA AND METHODOLOGY

Data used in this research were collected using the National Weather Service's Storm Prediction Center (SPC) Svrplot v2.0 program (Hart 2004), which lists all severe convective wind and severe hail events from 1955 to 2004, as well as all tornado events from 1950 to 2004. All severe weather events can be categorized by event type, intensity, occurrence time, and with tornadoes, according to F-scale, path length and width. Event times are referenced to Eastern Standard Time (EST).

The Fujita Scale is used to classify tornadoes into five categories based on the amount of damage that is produced (Fujita 1981). Estimated wind speeds are then provided for each category, with F0 being the weakest and F5 the strongest ([Table 1](#)). The total number of tornadoes included in this database is less than the number of tornado segments identified in this study. Many tornadoes track through more than one county. To accurately account for the number of tornado occurrences in a given county, these multi-county tornadoes are broken up into tornado segments. For example, a tornado that tracks uninterrupted through three counties would be identified as three separate events (three segments). Beginning in 1994, tornado occurrences were recorded according to this method. Thus, according to [McCarthy and Schaefer \(2004\)](#), this change in procedure has led to an increase in the number of tornado segments recorded since 1994.

A severe convective wind event, according to the National Weather Service, is defined by winds of 50 kts (58 mph) or greater, or winds causing damage to structures. Likewise, a severe hail event,

also defined by the National Weather Service, requires hail size to reach a minimum diameter of 0.75 inches.

3. COUNTY WARNING AREA DEMOGRAPHICS AND TOPOGRAPHY

3.1 Demographics

The Wilmington, Ohio, CWA covers over 20,000 square miles ([Fig. 2](#)), with a total population for the area estimated at nearly 5.5 million people as of 2000 ([Fig. 3](#)). There are two major metropolitan areas that account for nearly 80% of the total population. The largest population center is the Tri-State metropolis across southwest Ohio, northern Kentucky and southeast Indiana, with a population base of around 3 million people. The Tri-State metropolis consists of the cities of Cincinnati, Hamilton, and Covington and extends northward to encompass Dayton, Xenia, and Springfield. The greater Columbus metropolitan area, which includes the cities of Newark, Lancaster, and Delaware, has a population base of nearly 1.5 million people ([U.S. Census Bureau 2000](#)).

The remainder of the forecast area is largely rural and sparsely populated. This contributes to an average population density across the CWA of 256 people per square mile ([Fig. 4](#)), lower than the population densities in the metropolitan areas. Some skewing of observed severe weather towards the higher population centers is likely given the uneven population distribution of people across the CWA.

3.2 Topography

The topography of the Wilmington, Ohio CWA varies, ranging from plains across west-central Ohio and eastern Indiana to gently rolling to hilly terrain over southern Ohio and northern Kentucky. The Till Plains of western Ohio mark the eastern extent of the Midwestern Corn Belt, and are found across the northwest portion of the CWA. Terrain becomes gently rolling and dotted by hills further to the south, as the Great Miami, Little Miami and Scioto rivers carve valleys oriented north-south through the CWA. More rugged terrain is found across southern Ohio and northern Kentucky, in closer proximity to the Ohio River and on the western fringes of the Appalachian plateau. Elevations throughout the CWA range from 500 to 1000 feet above ground level (AGL), as shown in [Fig. 5](#).

4. SEVERE WEATHER CLIMATOLOGY

A total number of 6,131 severe weather events occurred across the Wilmington, Ohio, CWA during the period 1950-2004. Of this total, 4,175 severe convective wind events were reported, or nearly 68% of the total number of events ([Fig. 6](#)). A total of 1,470 severe hail events ([Fig. 7](#)) and 405 tornadoes ([Fig. 8](#)) were reported. As a whole, observed events tended to be more concentrated in the metropolitan areas surrounding Cincinnati, Columbus and Dayton. Fewer events, however, were

reported across more rural areas of the Wilmington CWA, in particular across southeast Indiana and portions of northern Kentucky and south-central Ohio.

4.1 Tornado Climatology

Between 1950 and 2004, 405 tornadoes, or 486 tornado segments (see Section 2) occurred across the Wilmington, Ohio, CWA. An annual distribution of tornadoes ([Fig. 9](#)) shows that despite a gradual increase in the number of reported tornadoes since the 1950s and 1960s, the number of tornadoes remains highly variable from year to year. This is due at least in part to the yearly variability of weather patterns that affect the Ohio Valley. The Palm Sunday outbreak in April 1965 and the Superoutbreak in April 1974 were two of the most significant tornado outbreaks to occur in recorded history across the Midwest and Ohio Valley. These tornado outbreaks were noteworthy in terms of the number of violent tornadoes (F4-F5) within each outbreak. 1992 was the most active year for tornadoes with 24. The number of tornadoes producing F2 damage and greater has largely remained unchanged since 1950. There has, however, been a noted increase in the yearly proportion of the total number of tornadoes to the number of significant tornadoes. This has likely been influenced by an overall increase in public severe weather awareness and the development and expansion of the trained spotter and amateur radio networks ([McCarthy and Schaefer 2004](#)).

Tornadoes were most commonly reported across the Wilmington, Ohio CWA between April and July ([Fig. 10](#)), making up approximately 73% of all tornadoes in this study. Many of the tornadoes that occurred during this same period (April-July) demonstrated the tendency to produce F2 damage or greater ([Fig. 11](#)). A minor secondary peak is noted in November, with half of these being F2 and greater. Tornadoes most frequently occurred between the mid-afternoon and evening hours ([Fig. 12](#)). Just over 50% of all tornadoes occurred between 3-7 PM EST, with the least amount of activity occurring between midnight and 10 AM EST.

While 80% of all tornado-related fatalities across the Wilmington CWA occurred during the late afternoon and evening hours, only 5% of the total number of fatalities took place between midnight and 1 PM ([Fig. 13](#)). Likewise, nearly 75% of all tornado-related injuries occurred during the late afternoon and evening, with 45% of these injuries occurring between 3-4 PM ([Fig. 14](#)). The anomalous spike during this time is clearly biased by the 3 April 1974 Xenia, Ohio, tornado, which accounted for 1,150 injuries alone.

Approximately 70% of all tornadoes examined across the Wilmington CWA between 1950 and 2004 were classified as “weak” tornadoes (F0-F1) with winds of 60-115 mph ([Fig. 15](#)). While close to half of all tornadoes were classified as F1, nearly 30% were categorized as significant tornadoes (F2-F5). The percentage of tornadoes then decreases by nearly a factor of three for each successive category. Violent tornadoes (F4-F5), while having made up only 4% of all tornadoes, accounted for approximately 82% of all tornado-related fatalities ([Fig. 16](#)) and 76% of all tornado-related injuries ([Fig. 17](#)). The 1974 Xenia, Ohio tornado, however, contributed significantly to the higher percentage of fatalities and injuries in the violent tornadoes category. This tornado, which was a part of the Superoutbreak, overwhelmingly produced the highest number of fatalities (36) and injuries (1150) of any tornado during the 1950-2004 period.

The tracks of all tornadoes producing F2 damage and greater across the Wilmington CWA can be seen in [Fig. 18](#). The map hints at a favored track pattern that extends from southern Indiana northeast through the Cincinnati area, then on into the Columbus area. A weaker signal can be seen across the far northern portion of the CWA, where tornadoes appear to track in a more west to east direction. This supports the findings of [Broyles and Crosbie \(2004\)](#), who provided evidence indicating that both of these favored tracks are actually portions of two smaller tornado alleys for long track tornadoes that produced F3 damage and greater over a 124-year period. They identified an area of increased long- track significant tornadoes extending from southern Missouri northeast into southwest Ohio, with a second area extending from central Illinois east-northeast into northwest Ohio.

To better assess path lengths and widths, tornadoes were grouped into three F-scale related categories (F0-F1, F2-F3, and F4-F5) and average path lengths and widths were calculated for each group. [Figure 19](#) indicates that the path length increases by nearly a factor of four for each category, with violent tornadoes rated as F4 and F5 averaging just over 32 miles in length. [Figure 20](#) shows a similar increase, but for path width, expanding by nearly a factor of three for each successive category.

4.2 Convective Wind Climatology

The annual distribution of severe convective wind events shows a dramatic increase in observed events over the period of study ([Fig. 21](#)). This corresponds to the national trend which has shown more than a 400% increase in the number of annual severe convective wind reports between 1970-2000 ([Weiss et al. 1999](#)). As with tornado reports, the development of the trained spotter and amateur radio networks have certainly contributed to this. Note that wind data are missing from 1972. This coincides with a change in procedure in the collection of wind data in the NOAA publication *Storm Data*. Prior to 1972, convective wind and hail data were collected by the U.S. Air Force ([Schaefer and Edwards 1999](#)). An increase in reports occurred in the 1980s, corresponding to the implementation of a national warning verification program within the National Weather Service.

An additional rise in annual reports in the early 1990s coincides with the deployment of the WSR-88D NEXRAD Radar network ([Weiss et al. 1999](#)).

Severe convective wind events have occurred in each month of the year across the Wilmington CWA. The number of events per month increase through the spring, and peak during June and July ([Fig. 22](#)). This peak in convective wind events corresponds with the annual peak in severe weather events during the early summer. The frequency of convective wind reports decrease during the late summer, with a secondary maximum observed in events noted in November. Damaging wind events are most frequent during the late afternoon and evening hours, with 67% of the total number of events occurring between 2-9 PM, and 52% between 3-7 PM ([Fig. 23](#)). This coincides with the time period when surface heating and atmospheric instability values are normally greatest. The frequency of severe convective wind events decreased during the late evening and overnight hours, with the minimum occurring between 4-10 AM.

4.3 Hail Climatology

The annual distribution of severe hail events shows a noted increase beginning in the early 1990s ([Fig. 24](#)). This distribution is similar to that previously noted for the severe convective wind events between 1955 and 2004 ([Fig. 21](#)), and corresponds to the overall increase in reports nationally since 1955. This too is likely influenced heavily by the increase in trained spotter and amateur radio networks (Schaefer et al. 2004). Fluctuations in synoptic patterns, particularly those supporting severe convection, will have an obvious influence on the total number of severe weather reports. This may be part of the reason behind both the 1997 and 1999 datasets indicating a significantly lower number of severe hail events when compared to the average annual number of hail events during the period 1994-2004.

For the period 1955-2004, severe hail was recorded in every month but December ([Fig. 25](#)). The frequency of severe hail events is greatest from April through June. As with the monthly tornado and severe convective wind distributions, a small secondary maximum is noted in November. Severe hail events are most frequent during the afternoon and early evening hours, with 65% of the total number of events occurring between 2-8 PM ([Fig. 26](#)). A minimum number of severe hail events occurred during the overnight and early morning hours, with only about 7% of the total number of events taking place between the hours of 1-11 AM.

Hail of 1.00 inch in diameter or smaller accounted for nearly 73% of the total number of severe hail events, with penny-size hail (0.75 inches in diameter) being the most widely reported hail size. Two and one-half (2.5) inch diameter hail and larger were rare in the CWA, making up only about 3% of the total number of severe hail reports ([Fig. 27](#)).

5. SUMMARY

Over 6,000 severe weather events consisting of tornadoes, severe convective wind and hail were compiled during the period 1950-2004 using the Svrplot v2.0 program from the National Weather Service's Storm Prediction Center. Events were categorized by severe weather type on annual, monthly and hourly time scales.

Tornadoes were found to be most frequent during the spring and early summer, primarily between April and July. Significant tornadoes rated as F2 and greater ([see Table 1](#)) peaked in April, and were strongly influenced by the large tornado outbreaks that occurred on 11 April 1965 and 3 April 1974. Severe convective wind events were most frequent between April and August, while severe hail events were most common between April and July. All three severe weather types showed a weaker secondary maximum in November. Additional research is needed to help understand this minor secondary peak.

The annual number of severe weather reports has increased dramatically since 1990, with nearly 70% of all of the reports included in this study occurring from that point. Several reasons can be attributed to this increase.

These include:

- 1) The development of the trained spotter and amateur radio networks beginning in the 1990s;
- 2) An increase in public severe weather awareness via the NWS, media and other sources;
- 3) Increases in population throughout the CWA;
- 4) Improved detection capabilities as a result of the advent and deployment of the WSR-88D Doppler radar network throughout the NWS in the early 1990s.

The information presented in this study identifies historical trends in location and types of severe weather at the Weather Forecast Office (WFO) in Wilmington, Ohio. The data presented within this climatology will provide forecasters with a background for evaluating the frequency and impact of different types of severe weather on temporal and spatial scales, and act as an additional resource for forecasters in anticipation of severe weather within the CWA. Also, it is envisioned that the results documented in this study will spawn subsequent locally focused severe convection climatologies that impact the Wilmington CWA.

ACKNOWLEDGMENTS

The author would like to thank John DiStefano (Science and Operations Officer) and Stephen Hrebenach (Senior Forecaster) at the WFO in Wilmington, Ohio, for their reviews and suggestions in association with this study.

REFERENCES

Broyles, J.C., and K.C. Crosbie, 2004: Evidence of smaller tornado alleys across the United States based on a long track F3-F5 tornado climatology study from 1880-2003. Preprints, *22nd Conf. Severe Local Storms*, Hyannis, MA, Amer. Meteor. Soc., P5.6.

Fujita, T.T., 1981: Tornadoes and downbursts in the context of generalized planetary scales. *J. Atmos. Sci.*, **38**, 1511-34.

Hart, J.A., 2004: SeverePlot v2, NCEP Storm Prediction Center, Norman, OK [Available online from: <http://www.spc.noaa.gov/software/svrplot2/>].

McCarthy, D.W., and J.T. Schaefer, 2004: Tornado trends over the past 30 years. Preprints, *14th Conf. Applied Climatology*, Seattle, WA, Amer. Meteor. Soc., 3.4.

Schaefer, J.T., J.J. Levit, S.J. Weiss and D.W. McCarthy, 2004: The Frequency of Large Hail over the Contiguous United States. Preprints, *14th Conf. Applied Climatology*, Seattle, WA, Amer. Meteor. Soc., 3.3.

Schaefer, J.T., and R. Edwards, 1999: The SPC tornado/severe thunderstorm database. Preprints, *11th Conf. Applied Climatology*, Dallas, TX, Amer. Meteor. Soc., 215-220.

United States Census Bureau. 2000 Census Statistics. [Available online from <http://www.census.gov/main/www/cen2000.html>], April, 2006.

Weiss, S.J., J.A. Hart and P.R. Janish, 2002: An examination of severe thunderstorm wind report climatology: 1970-1999. Preprints, *21st Conf. Severe Local Storms*, San Antonio, TX, Amer. Meteor. Soc., 446-449.

Table 1. The Fujita Damage Scale (Fujita, 1981)

| <u>Scale</u> | <u>Wind Speed (mph)</u> | <u>Tornado Character</u> | <u>Damage Intensity</u> |
|---------------------|--------------------------------|---------------------------------|--------------------------------|
| F0 | 40-72 | Weak | Light |
| F1 | 73-112 | Weak | Moderate |
| F2 | 113-157 | Strong | Considerable |
| F3 | 158-206 | Strong | Severe |
| F4 | 207-260 | Violent | Devastating |
| F5 | 261-318 | Violent | Incredible |

FIGURES




Figure 1. The 52 counties that comprise the Wilmington Ohio County Warning Area (CWA), including southwest Ohio, north-central Kentucky and southeast Indiana. The blue line indicates the state borders.




Figure 2. The total area in square miles by county across the Wilmington, Ohio CWA.




Figure 3. The population by county across the Wilmington, Ohio CWA based on the 2000 U.S. Census Bureau numbers.




Figure 4. The county population density across the Wilmington, Ohio CWA based on the 2000 U.S. Census Bureau numbers.




Figure 5. A high-resolution topographic map of the Wilmington, Ohio CWA, taken from the Advanced Weather Information Processing System (AWIPS). Heights range from 500 feet (lighter yellow-orange) to 1500 feet AGL (purple).




Figure 6. The total number of severe convective wind events across the Wilmington, Ohio, CWA from 1955-2004.




Figure 7. The total number of severe hail events across the Wilmington, Ohio CWA from 1950-2004.




Figure 8. The total number of tornadoes across the Wilmington, Ohio CWA from 1950-2004.




Figure 9. The annual distribution of tornadoes across the Wilmington, Ohio CWA, from 1950-2004.




Figure 10. The monthly distribution of tornadoes across the Wilmington, Ohio CWA from 1950-2004.




Figure 11. The monthly distribution of significant tornadoes (F2-F5) across the Wilmington, Ohio, CWA from 1950-2004.




Figure 12. The hourly distribution of tornadoes across the Wilmington, Ohio CWA from 1950-2004. Hours are in Eastern Standard Time.




Figure 13. Tornado-related fatalities by hour across the Wilmington, Ohio CWA from 1950-2004. Hours are in Eastern Standard Time.




Figure 14. Tornado-related injuries by hour across the Wilmington, Ohio CWA from 1950-2004. Hours are in Eastern Standard Time.




Figure 15. Percentage of tornadoes by F-Scale across the Wilmington, Ohio CWA from 1950-2004.




Figure 16. The number of tornado fatalities by F-scale across the Wilmington, Ohio CWA from 1950 to 2004.




Figure 17. The number of tornado injuries by F-scale across the Wilmington, Ohio CWA from 1950 to 2004.




Figure 18. The significant tornado (F2-F5) tracks across the Wilmington, Ohio CWA from 1950-2004.




Figure 19. The average tornado path length (in miles) for weak (F0-F1), strong (F2-F3) and violent (F4-F5) tornadoes across the Wilmington, Ohio CWA from 1950-2004.




Figure 20. The average tornado width (in yards) for weak (F0-F1), strong (F2-F3) and violent (F4-F5) tornadoes across the Wilmington, Ohio CWA from 1950-2004.




Figure 21. The annual distribution of severe convective wind events across the Wilmington, Ohio, CWA from 1955-2004. Wind data from 1972 are missing.




Figure 22. The monthly distribution of severe convective wind events across the Wilmington, Ohio CWA from 1955-2004.




Figure 23. The hourly distribution of severe convective wind events across the Wilmington, Ohio, CWA from 1955-2004. Hours are in Eastern Standard Time.




Figure 24. The annual distribution of severe hail events across the Wilmington, Ohio CWA from 1955-2004.




Figure 25. The monthly distribution of severe hail events across the Wilmington, Ohio CWA from 1955-2004.




Figure 26. The hourly distribution of severe hail events across the Wilmington, Ohio CWA.




Figure 27. The distribution of severe hail by size across the Wilmington, Ohio CWA.

NWS ER 46 An Objective Method of Forecasting Summertime Thunderstorms. John F. Townsend and Russell J. Younkin. May 1972. (COM-72-10765).

NWS ER 47 An Objective Method of Preparing Cloud Cover Forecasts. James R. Sims. August 1972. (COM-72-11382).

NWS ER 48 Accuracy of Automated Temperature Forecasts for Philadelphia as Related to Sky Condition and Wind Direction. Robert B. Wassall. September 1972. (COM-72-11473).

NWS ER 49 A Procedure for Improving National Meteorological Center Objective Precipitation Forecasts. Joseph A. Ronco, Jr. November 1972. (COM-73-10132).

NWS ER 50 PEATMOS Probability of Precipitation Forecasts as an Aid in Predicting Precipitation Amounts. Stanley E. Wasserman. December 1972. (COM-73-10243).

NWS ER 51 Frequency and Intensity of Freezing Rain/Drizzle in Ohio. Marvin E. Miller. February 1973. (COM-73-10570).

NWS ER 52 Forecast and Warning Utilization of Radar Remote Facsimile Data. Robert E. Hamilton. July 1973. (COM-73-11275).

NWS ER 53 Summary of 1969 and 1970 Public Severe Thunderstorm and Tornado Watches Within the National Weather Service, Eastern Region. Marvin E. Miller and Lewis H. Ramey. October 1973. (COM-74-10160).

NWS ER 54 A Procedure for Improving National Meteorological Center Objective Precipitation Forecasts - Winter Season. Joseph A. Ronco, Jr. November 1973. (COM-74-10200).

NWS ER 55 Cause and Prediction of Beach Erosion. Stanley E. Wasserman and David B. Gilhousen. December 1973. (COM-74-10036).

NWS ER 56 Biometeorological Factors Affecting the Development and Spread of Planet Diseases. V.J. Valli. July 1974. (COM-74-11625/AS).

NWS ER 57 Heavy Fall and Winter Rain In The Carolina Mountains. David B. Gilhousen. October 1974. (COM-74-11761/AS).

NWS ER 58 An Analysis of Forecasters' Propensities In Maximum/Minimum Temperature Forecasts. I. Randy Racer. November 1974. (COM-75-10063/AS).

NWS ER 59 Digital Radar Data and its Application in Flash Flood Potential. David D. Sisk. March 1975. (COM-75-10582/AS).

NWS ER 60 Use of Radar Information in Determining Flash Flood Potential. Stanley E. Wasserman. December 1975. (PB250071/AS).

NWS ER 61 Improving Short-Range Precipitation Guidance During the Summer Months. David B. Gilhousen. March 1976. (PB256427).

NWS ER 62 Locally Heavy Snow Downwind from Cooling Towers. Reese E. Otts. December 1976. (PB263390/AS).

NWS ER 63 Snow in West Virginia. Marvin E. Miller. January 1977. (PB265419/AS).

NWS ER 64 Wind Forecasting for the Monongahela National Forest. Donald E. Risher. August 1977. (PB272138/AS).

NWS ER 65 A Procedure for Spraying Spruce Budworms in Maine during Stable Wind Conditions. Monte Glovinsky. May 1980. (PB80-203243).

NWS ER 66 Contributing Factors to the 1980-81 Water Supply Drought, Northeast U.S. Solomon G. Summer. June 1981. (PB82-172974).

NWS ER 67 A Computer Calculation and Display System for SLOSH Hurricane Surge Model Data. John F. Townsend. May 1984. (PB84-198753).

NWS ER 68 A Comparison Among Various Thermodynamic Parameters for the Prediction of Convective Activity. Hugh M. Stone. April 1985. (PB85-206217/AS).

NWS ER 69 A Comparison Among Various Thermodynamic Parameters for the Prediction of Convective Activity, Part II. Hugh M. Stone. December 1985. (PB86-142353/AS).

NWS ER 70 Hurricane Gloria's Potential Storm Surge. Anthony G. Gigi and David A. Wert. July 1986. (PB86-226644/AS).

NWS ER 71 Washington Metropolitan Wind Study 1981-1986. Clarence Burke, Jr. and Carl C. Ewald. February 1987. (PB87-151908/AS).

NWS ER 72 Mesoscale Forecasting Topics. Hugh M. Stone. March 1987. (PB87-180246/AS).

NWS ER 73 A Procedure for Improving First Period Model Output Statistics Precipitation Forecasts. Antonio J. Lacroix and Joseph A. Ronco, Jr. April 1987. (PB87-180238/AS).

NWS ER 74 The Climatology of Lake Erie's South Shoreline. John Kwiatkowski. June 1987. (PB87-205514/AS).

NWS ER 75 Wind Shear as a Predictor of Severe Weather for the Eastern United States. Hugh M. Stone. January 1988. (PB88-157144).

NWS ER 76 Is There A Temperature Relationship Between Autumn and the Following Winter? Anthony Gigi. February 1988. (PB88-173224).

NWS ER 77 River Stage Data for South Carolina. Clara Cillentine. April 1988. (PB88-201991/AS).

NWS ER 78 National Weather Service Philadelphia Forecast Office 1987 NOAA Weather Radio Survey & Questionnaire. Robert P. Wanton. October 1988. (PB89-111785/AS).

NWS ER 79 An Examination of NGM Low Level Temperature. Joseph A. Ronco, Jr. November 1988. (PB89-122543/AS).

NWS ER 80 Relationship of Wind Shear, Buoyancy, and Radar Tops to Severe Weather 1988. Hugh M. Stone. November 1988. (PB89-1222419/AS).

NWS ER 81 Relation of Wind Field and Buoyancy to Rainfall Inferred from Radar. Hugh M. Stone. April 1989. (PB89-208326/AS).

NWS ER 82 Second National Winter Weather Workshop, 26-30 Sept. 1988: Postprints. Laurence G. Lee. June 1989. (PB90-147414/AS).

NWS ER 83 A Historical Account of Tropical Cyclones that Have Impacted North Carolina Since 1586. James D. Stevenson. July 1990. (PB90-259201).

NWS ER 84 A Seasonal Analysis of the Performance of the Probability of Precipitation Type Guidance System. George J. Maglaras and Barry S. Goldsmith. September 1990. (PB93-160802).

NWS ER 85 The Use of ADAP to Examine Warm and Quasi-Stationary Frontal Events in the Northeastern United States. David R. Vallee. July 1991. (PB91-225037).

NWS ER 86 Rhode Island Hurricanes and Tropical Storms A Fifty-Six Year Summary 1936-1991. David R. Vallee. March 1993. (PB93-162006).

NWS ER 87 Post-print Volume, Third National Heavy Precipitation Workshop, 16-20 Nov. 1992. April 1993. (PB93-186625).

NWS ER 88 A Synoptic and Mesoscale Examination of the Northern New England Winter Storm of 29-30 January 1990. Robert A. Marine and Steven J. Capriola. July 1994. (PB94-209426).

NWS ER 89 An Initial Comparison of Manual and Automated Surface Observing System Observations at the Atlantic City, New Jersey, International Airport. James C. Hayes and Stephan C. Kuhl. January 1995.

NWS ER 90 Numerical Simulation Studies of the Mesoscale Environment Conducive to the Raleigh Tornado. Michael L. Kaplan, Robert A. Rozumalski, Ronald P. Weglarz, Yuh-Lang Lin, Steven Businger, and Rodney F. Gonski. November 1995.

NWS ER 91 A Climatology of Non-convective High Wind Events in Western New York State. Thomas A. Niziol and Thomas J. Paone. April 2000.

NWS ER 92 Tropical Cyclones Affecting North Carolina Since 1586 - An Historical Perspective. James E. Hudgins. April 2000.

NWS ER 93 A Severe Weather Climatology for the Wilmington, NC WFO County Warning Area. Carl R., Morgan. October 2001.

NWS ER 94 Surface-based Rain, Wind, and Pressure Fields in Tropical Cyclones over North Carolina since 1989. Joel Cline. June 2002.

NWS ER 95 A Severe Weather Climatology for the Charleston, South Carolina, WFO County Warning Area. Stephen Brueske, Lauren Plourd, Matthen Volkmer. July 2002.

NWS ER 96 A Severe Weather Climatology for the WFO Wakefield, VA County Warning Area. Brian T. Cullen. May 2003. (PB2003-105462).

NWS ER 97 Severe Weather Climatology for the Columbia, SC WFO County Warning Area. Leonard C. Vaughan. September 2003. (PB2004-100999).

NWS ER 98 Climatology of Heavy Rainfall Associated with Tropical Cyclones Affecting the Central Appalachians. James Hudgins, Steve Keighton, Kenneth Kostura, Jan Jackson. September 2005.

NWS ER 99 A Severe Weather Climatology for the WFO Blacksburg, Virginia, County Warning Area. Robert Stonefield, James Hudgins. January 2007.

NWS ER 100 Tropical Cyclones Affecting North Carolina Since 1586 - An Historical Perspective. James E. Hudgins. October 2007.

NWS ER 101 A Severe Weather Climatology for the Raleigh, NC County Warning Area. Clyde Brandon Locklear. May 2008.

NOAA SCIENTIFIC AND TECHNICAL PUBLICATIONS

The National Oceanic and Atmospheric Administration was established as part of the Department of Commerce on October 3, 1970. The mission responsibilities of NOAA are to assess the socioeconomic impact of natural and technological changes in the environment and to monitor and predict the state of the solid Earth, the oceans and their living resources, the atmosphere, and the space environment of the Earth.

The major components of NOAA regularly produce various types of scientific and technical information in the following kinds of publications:

PROFESSIONAL PAPERS--Important definitive research results, major techniques, and special investigations.

CONTRACT AND GRANT REPORTS--Reports prepared by contractors or grantees under NOAA sponsorship.

ATLAS--Presentation of analyzed data generally in the form of maps showing distribution of rainfall, chemical and physical conditions of oceans and atmosphere, distribution of fishes and marine mammals, ionospheric conditions, etc.

TECHNICAL SERVICE PUBLICATIONS--Reports containing data, observations, instructions, etc. A partial listing includes data serials; prediction and outlook periodicals; technical manuals, training papers, planning reports, and information serials; and miscellaneous technical publications.

TECHNICAL REPORTS--Journal quality with extensive details, mathematical developments, or data listings.

TECHNICAL MEMORANDUMS--Reports of preliminary, partial, or negative research or technology results, interim instructions, and the like.



Information on availability of NOAA publications can be obtained from:

NATIONAL TECHNICAL INFORMATION SERVICE
U.S. DEPARTMENT OF COMMERCE
5285 PORT ROYAL ROAD
SPRINGFIELD, VA 22161